

# DNP leader role modeling the use of evidence to decrease catheter associated urinary tract infections with leader rounding using a checklist Elizabeth Menschner, DNP, MAS, MSN, RN, NEA-BC

Temple University Hospital 3401 N Broad St. Phila., Pa. 215-707-9491 Elizabeth.Menschner@tuhs.temple.edu

# **PICO QUESTION**

- ${\bf P}$  In a dult medical and neurosurgical intensive care patients with ind welling urinary catheters
- I What is the effect of DNP leader role modeling daily nurse leader rounding and the use of a criteria-based reminder sheet
- C Compared with no reminder sheet
- $\mathbf{O}-\mathrm{On}$  indwelling catheter days and the rate of catheter acquired urinary tract infections
- T Over a six-week timeframe

### **ABSTRACT**

### Background:


It has been estimated that each year, more than 13,000 deaths are associated with CAUTI's (Magill et al., 2014). Decreasing use of indwelling catheters through removal when no longer clinically indicated is the key to decreasing or eliminating CAUTI. Approximately 12%-16% of adult hospitalized inpatients will have an indwelling urinary catheter at some time during their inpatient admission. Each day the indwelling urinary catheter remains, a patient has a 3% to 7% increased risk of acquiring a CAUTI (Burton, Edwards, Srinivasan, Fridkin, & Gould, 2011).


### Purpose:

The purpose of this project was to determine if nurse leader rounding using a checklist to prompt removal when indwelling urinary catheters are no longer clinically indicated would decrease catheter days and CAUTI infections compared to nurse leaders not rounding.

### **METHODS**

- •A process improvement plan was drafted
  - Utilized Focus Plan-Do-Check -Act framework (PDCA)
  - Preventing Catheter-Associated Urinary Tract Infections
  - · Four Components of Care:
  - · Avoid unnecessary urinary catheters
  - · Insert urinary catheters using aseptic technique
  - · Maintain urinary catheters based on recommended guidelines
  - Daily leader rounding using a checklist for urinary catheter necessity and removal





## **LIMITATIONS**

- One limitation to the study included the fact that it was a single site study in a multisystem organization.
- Another limitation was that the study had a short duration of a six week period of time. It may be beneficial to evaluate a longer period of time
- There were two CAUTIs during the study period. One CAUTI was attributed to the neurosurgical intensive care unit that occurred when the patient was in another location within the hospital.
- Another limitation included the fact that there are many other elements that contribute to CAUTI and length of time the catheter is in place is not the only rate limiting factor.

# RECOMMENDATIONS FOR PRACTICE

- · CAUTI infections should not occur in the hospitalized patient.
- · CAUTI is considered a "never event" by TJC.
- CAUTI infections impact the quality of life for our patients and are considered a nurse sensitive indicator.
- DNP leader's to role model use evidence based practice through daily review of the ongoing need for indwelling urinary catheters supports empowering the clinical nurse to remove catheters following the nurse driven protocol.
- Outcomes of this study was shared with the study site and system CAUTI committee

# REFERENCES

AHRQ. (2015). http://www.ahrq.gov/health-care-information/topics/topic-hai.html

Alexaitis, I., & Broome, B., (2014). Implementation of a nurse-driven protocol to prevent catheter-associated urinary track infections. *Journal of Nursing Care Quality*, 29(3), 245-252. Doi:10.1097/NCQ.0000000000000001

Bader, M. K., Palmer, S., Stalcup, C., & Shaver, T. (2003). Using a FOCUS-PDCA quality improvement model for applying the severe traumatic brain injury guidelines to practice: Process and outcomes. Evidence - Based Nursing, 6(1), 6. doi:http://dx.doi.org.ezproxy2.library.drexl.edu/10.1136/ebn.61.6

Burton, D. C., Edwards, J. R., Srinivasan, A., Fridkin, S. K., & Gould, C. V. (2011, August). Trends in Catheter-Associated Urinary Tract Infections in Adult Intensive Care Units-United States, 190-2007. Infection Control and Hospital Epidemiology, 32(8), 748-756. <a href="http://dx.doi.org/23.235.32.0">http://dx.doi.org/23.235.32.0</a>

Centers for Disease Control and Prevention. (2012), www.cdc.gov/infection/

Chen, Y., Chi, M., Chen, Y., Chan, Y., Chou, S., & Wang, F. (2013). Using a criteria-based reminder to reduce use of indwelling urinary catheters and decrease urinary tract infections. AJCC, 22(2), 105-114. http://dx.doi.org/http://dx.doi.org/10.0437/iacg.2013464

Parry, M. F., Grant, B., & Sestovic, M. (2013). Successful reduction in catheter-associated urinary tract infections: Focus on nurse-directed catheter removal.

American Journal of Infection Control, 41(), 1178-1181. http://dx.doi.org/10.1016/j.ajic.2013.03.296